

THE AUSLANDER AND RINGEL-TACHIKAWA THEOREM FOR SUBMODULE EMBEDDINGS

AUDREY MOORE

ABSTRACT. Auslander and Ringel-Tachikawa have shown that for an artinian ring R of finite representation type, every R -module is the direct sum of finitely generated indecomposable R -modules. In this paper, we will adapt this result to finite representation type full subcategories of the module category of an artinian ring which are closed under subobjects and direct sums and contain all projective modules. In particular, the results in this paper hold for subspace representations of a poset, in case this subcategory is of finite representation type.

1. INTRODUCTION

Auslander, [4, Corollary 4.8], Ringel and Tachikawa [12, Corollary 4.4] showed that if R is an artinian ring of finite representation type, then every R -module is the sum of finitely generated indecomposable R -modules. In this paper, we are interested in a relative version of this result: Let $\text{Mod } R$ be a module category which may not be of

⁰2000 MSC. Primary: 16G60, Secondary: 16G70, 20E15, 47A15

⁰This research was partially supported by NSF DDEP grant #0831369

finite type, and $\mathcal{S} \subseteq \text{Mod } R$ a subcategory which has only finitely many finite length indecomposable objects up to isomorphism. Under which circumstances is every object in $\mathcal{S}$ a direct sum of finite length indecomposable objects?

Theorem 1. *Let R be an artinian ring, $\mathcal{S}$ a full subcategory of $\text{Mod } R$ which is closed under direct sums and subobjects with $R_R \in \mathcal{S}$. If there are only finitely many finite length indecomposable objects in $\mathcal{S}$, up to isomorphism, then every object in $\mathcal{S}$ is a direct sum of finitely generated indecomposable subobjects. In particular, every indecomposable module in $\mathcal{S}$ has finite length.*

The proof of Theorem 1 will be given in Section 4, where we will adapt a proof of the classical Auslander and Ringel-Tachikawa Theorem, which was given in lecture notes by W. Zimmermann [14]. As representations of posets are a motivating example, we will start with a slightly more restricted situation in Section 2, and use Auslander-Reiten theory to prove Proposition 2 below, a restatement of Theorem 1 in this situation in terms of Auslander-Reiten theory. The benefit of this method of proof is that we obtain not only the result, but also

information about the individual indecomposables and the almost split morphisms.

Proposition 2. *Let R be a right Morita ring and $\mathcal{S}$ a full subcategory of $\text{Mod } R$ which is closed under direct sums and summands. If Γ is a finite component of the Auslander-Reiten quiver for $\mathfrak{s} = \mathcal{S} \cap \text{mod } R$ such that Γ contains a projective generator, only endofinite modules, and each Auslander-Reiten sequence in Γ is an Auslander-Reiten sequence in $\mathcal{S}$, then $\mathfrak{s}$ has Auslander-Reiten sequences in $\mathcal{S}$, Γ is the Auslander-Reiten quiver for $\mathfrak{s}$ and every object in $\mathcal{S}$ is a direct sum of indecomposable modules in Γ .*

Section 3 will focus on representations of posets with coefficients in a right Morita ring, illustrating the results from Section 2. As an example we obtain the following application to invariant subspaces of linear operators. This situation is of particular interest since the category of modules over the incidence algebra has infinite type, and hence does not satisfy the conditions of the classical version of the Auslander and Ringel-Tachikawa Theorem.

Proposition 3. *Let k be any field, V a possibly infinite dimensional k -vector space, $T : V \rightarrow V$ a k -linear operator which acts on V with*

nilpotency index 2, and let V_1, V_2, V_3 be subspaces of V which are invariant under the action of T and such that $V_1 \subseteq (V_2 \cap V_3)$. Then V has a direct sum decomposition $V = \bigoplus_i W_i$ into T -invariant vector spaces such that $V_j = \bigoplus_i V_j \cap W_i$ holds for $j = 1, 2, 3$ and where each W_i is isomorphic to exactly one of the 25 systems pictured in Section 3.

Related Results: For P the one point poset and $\Lambda = \frac{\mathbf{Z}}{p^n}$ we are dealing with the problem posed by Birkhoff in 1934 to classify all subgroups of any finite abelian p^n -bounded group B , up to an automorphism of B . For $n \leq 5$, Richman and Walker have shown that any subspace representation of P is the direct sum of finitely generated indecomposable subspace representations of P [8]. This is particularly interesting in the case $n = 4$ or $n = 5$, where the module category of the incidence algebra is not of finite representation type. Recently, C.M. Ringel [9] has given a proof of the Auslander and Ringel-Tachikawa Theorem in the case where R is an artin algebra which uses the Gabriel-Roiter measure and yields a method for splitting off direct summands.

Acknowledgements: This paper is part of the author's Ph.D. thesis advised by Markus Schmidmeier at Florida Atlantic University. The author would like to thank the representation theory group in Bielefeld, where this paper was completed, for their hospitality and the

NSF for supporting this international exchange through DDEP grant #0831369.

2. AUSLANDER-REITEN SEQUENCES AND A RESULT ABOUT INDECOMPOSABLES

In this section, let R be a (two-sided) artinian ring, $\text{Mod } R$ the category of right R -modules, $\mathcal{S}$ a full subcategory of $\text{Mod } R$ which is closed under direct sums and summands with $R_R \in \mathcal{S}$, and $\mathfrak{s} = \mathcal{S} \cap \text{mod } R$, the full subcategory of $\mathcal{S}$ which has as objects the finitely generated modules in $\mathcal{S}$. We will give sufficient conditions for $\mathfrak{s}$ to have Auslander-Reiten sequences with factorization property in $\mathcal{S}$ and deduce that if $\mathfrak{s}$ is of finite type and has endofinite indecomposables, then any object in $\mathcal{S}$ can be written as a direct sum of finite length indecomposable modules. The terminology below regarding Auslander-Reiten sequences is adapted from [5].

Let $B, C \in \mathfrak{s}$. A morphism $g : B \rightarrow C$ is *right almost split in $\mathcal{S}$* if g is not a split epimorphism, and for any $h : X \rightarrow C$ which is not a split epimorphism, with $X \in \mathcal{S}$, the map h factors through g . Define left almost split in $\mathcal{S}$ dually. Then we say $\mathfrak{s}$ *has right almost split morphisms in $\mathcal{S}$* if for any indecomposable object $C \in \mathfrak{s}$, there is an object $B \in \mathfrak{s}$

and an $f : B \rightarrow C$ which is right almost split in $\mathcal{S}$. Define $\mathfrak{s}$ has left almost split morphisms in $\mathcal{S}$ dually. We say $\mathfrak{s}$ is functorially finite in $\text{mod } R$ with respect to $\mathcal{S}$ if $\mathfrak{s}$ is functorially finite in $\text{mod } R$, and the approximations have the factorization property for test objects in $\mathcal{S}$.

A non-split exact sequence $0 \rightarrow A \xrightarrow{f} B \xrightarrow{g} C \rightarrow 0$ with $A, B, C \in \mathfrak{s}$ is an *Auslander-Reiten sequence* in $\mathcal{S}$ if f is left almost split in $\mathcal{S}$ and g is right almost split in $\mathcal{S}$. We say $\mathfrak{s}$ has *Auslander-Reiten sequences* in $\mathcal{S}$ if the following conditions hold:

- (i) The category $\mathfrak{s}$ has left and right almost split morphisms in $\mathcal{S}$.
- (ii) For each indecomposable, non-projective $C \in \mathfrak{s}$, there is an Auslander-Reiten sequence in $\mathcal{S}$ ending in C with objects in $\mathfrak{s}$.
- (iii) For each indecomposable, non-injective $A \in \mathfrak{s}$, there is an Auslander-Reiten sequence in $\mathcal{S}$ beginning in A with objects in $\mathfrak{s}$.

Proposition 4. *Let $\mathcal{S}$ be a full subcategory of $\text{Mod } R$ which is closed under direct sums and summands such that $\mathfrak{s} = \mathcal{S} \cap \text{mod } R$ is functorially finite in $\text{mod } R$ with respect to $\mathcal{S}$. If $\text{mod } R$ has Auslander-Reiten sequences in $\text{Mod } R$, then $\mathfrak{s}$ has Auslander-Reiten sequences in $\mathcal{S}$.*

Proof. From Auslander and Smalø, we know that since $\mathfrak{s}$ is functorially finite in $\text{mod } R$, $\mathfrak{s}$ has Auslander-Reiten sequences in $\mathfrak{s}$ [5, section 4].

We are now ready to verify requirements (i) through (iii).

(i): Let $A \in \mathfrak{s}$ be indecomposable. Since $A \in \text{mod } R$, there is a left almost split morphism $f : A \rightarrow B$ in $\text{Mod } R$ with $B \in \text{mod } R$. Let $b : B \rightarrow B'$ be a left approximation of B in $\mathfrak{s}$. Then the composition $b \circ f : A \rightarrow B'$ is left almost split in $\mathcal{S}$. A dual construction yields right almost split morphisms in $\mathcal{S}$.

(ii): Let $C \in \mathfrak{s}$ be an indecomposable non-projective module. Then from (i), there is a right almost split morphism $g : B \rightarrow C$ in $\mathcal{S}$ with $B \in \mathfrak{s}$. Since B is a finite length module, we get a minimal version of this morphism by decomposing $B = B' \oplus B''$, with $g|_{B'} : B' \rightarrow C$ right minimal and $g|_{B''} = 0$ [6, I Theorem 2.2]. So $g|_{B'}$ is a minimal right almost split morphism, and we have an exact sequence

$$\mathcal{E} : 0 \rightarrow A \xrightarrow{f} B \xrightarrow{g|_{B'}} C \rightarrow 0$$

Consider the Auslander-Reiten sequence in $\mathfrak{s}$ ending in C : $0 \rightarrow \tilde{A} \xrightarrow{\tilde{f}} \tilde{B} \xrightarrow{\tilde{g}} C \rightarrow 0$. Since the minimal right almost split morphism in $\mathfrak{s}$ ending in C is unique up to isomorphism, we get an isomorphism h and a corresponding kernel map k such that the diagram commutes [6,

V Proposition 1.4]:

$$\begin{array}{ccccccc}
 0 & \longrightarrow & A & \xrightarrow{f} & B' & \xrightarrow{g|_{B'}} & C \longrightarrow 0 \\
 & & \downarrow k & & \downarrow h & & \parallel \\
 0 & \longrightarrow & \tilde{A} & \xrightarrow{\tilde{f}} & \tilde{B} & \xrightarrow{\tilde{g}} & C \longrightarrow 0
 \end{array}$$

Hence k is an isomorphism, so we see that $A \in \mathfrak{s}$ and $\mathcal{E}$ is an Auslander-Reiten sequence in $\mathfrak{s}$. Since $\mathfrak{s}$ has left almost split morphisms in $\mathcal{S}$, there is a left minimal almost split morphism $\hat{f} : A \rightarrow \hat{B}$ in $\mathcal{S}$ with $\hat{B} \in \mathfrak{s}$. Then $\hat{f}$ is also a minimal left almost split morphism in $\mathfrak{s}$, and so $\hat{f} \cong f$ by uniqueness of minimal almost split morphisms [6, V Proposition 1.5]. So f is minimal right almost split in $\mathcal{S}$, and $\mathcal{E}$ is an Auslander-Reiten sequence in $\mathcal{S}$.

(iii) Dual to the proof of (ii). $\square$

Notice that the assumption $\text{mod } R$ has Auslander-Reiten sequences in $\text{Mod } R$ must be stated in the previous proposition. This assumption is satisfied whenever R is an artin algebra [3, Theorem 3.9], but for R an artinian ring this may not be the case [13].

We say an additive subcategory of $\text{Mod } R$ is of *finite (representation) type* if it has only finitely many finite length indecomposable objects, up to isomorphism.

Corollary 5. *Let $\mathcal{S}$ be a full subcategory of $\text{Mod } R$ which is closed under direct sums and summands with $R_R \in \mathcal{S}$, and such that $\mathfrak{s} = \mathcal{S} \cap \text{mod } R$ is of finite type and has Auslander-Reiten sequences in $\mathcal{S}$. Then every non-zero object in $\mathcal{S}$ has a non-zero summand of finite length, and in particular each indecomposable in $\mathcal{S}$ has finite length.*

Proof. Let $X \in \mathcal{S}$ be non-zero. Then there is a nonzero morphism $f : P \rightarrow X$ with P an indecomposable projective. By assumption, $P \in \mathcal{S}$. If f is a split monomorphism, then X has a direct summand isomorphic to P , and we are done. So suppose f is not a split monomorphism. By assumption, $\mathfrak{s}$ has Auslander-Reiten sequences in $\mathcal{S}$, so there is a left almost split morphism starting at P , say $g_1 : P \rightarrow B_1$. Since f is not a split monomorphism, it factors over g_1 , so there is a map $f_1 : B_1 \rightarrow X$ such that $f = f_1 \circ g_1$. Also, since $B_1 \in \mathfrak{s}$, it is finitely generated, and we can write it as a sum of indecomposables: $B_1 = B_{1,1} \oplus B_{1,2} \oplus \dots \oplus B_{1,n}$, and the maps f_1 and g_1 are given by the maps on the indecomposable summands, say $f_1 = (f_{1,i})$, $g_1 = (g_{1,i})$. Since f is non-zero, f_1 is nonzero, and there is some i with $0 \neq f_{1,i} : B_{1,i} \rightarrow X$, say i_1 . Either f_{1,i_1} is a split monomorphism and we are done, or we continue.

Since $\mathfrak{s}$ is of finite type, there are only finitely many finitely generated indecomposables in $\mathfrak{s}$. Let m be the maximum length of these

indecomposable R -modules. By the Harada-Sai Lemma, this process must terminate after at most $2^m - 1$ steps, so we get a split monomorphism $B_{j,k} \rightarrow X$ for some indecomposable $B_{j,k} \in \mathfrak{s}$, and we see that X has a non-zero finite length direct summand. $\square$

Before proving the main result of the section, it may be useful to recall that a *Morita Ring* is an artinian ring such that the injective envelopes of simple right R -modules are finitely generated, and the notion of purity below.

Lemma 6. *The following are equivalent for a short exact sequence*

$$\mathcal{E} : 0 \rightarrow A \xrightarrow{f} B \xrightarrow{g} C \rightarrow 0$$

- i *For all ${}_R X$, $f \otimes 1_X$ is a monomorphism.*
- ii *For all finitely presented N_R , $\text{Hom}(N, g)$ is an epimorphism.*

If $\mathcal{E}$ satisfies these conditions, then $\mathcal{E}$ is called a *pure exact sequence*, and $f(A) \subseteq B$ is a pure submodule. So split exact sequences must be pure exact, and pure exact sequences are split exact if C is finitely presented.

We can now prove Proposition 2.

Proof. Since Γ is finite and contains a projective generator, all indecomposables in $\mathfrak{s}$ occur in Γ , so $\mathfrak{s}$ is of finite type. Since the almost split morphisms in Γ are almost split in $\mathcal{S}$, $\mathfrak{s}$ has Auslander-Reiten sequences in $\mathcal{S}$ and Γ is the Auslander-Reiten quiver for $\mathfrak{s}$. Let $X \in \mathcal{S}$, and $\mathcal{Z}$ be the set of all (internal) direct sums of finite length indecomposable direct summands of X . Then $\mathcal{Z}$ is non-empty and partially ordered by the direct sum relation. Furthermore, by Azumaya's Theorem [1, Theorem 12.6] each chain in $\mathcal{Z}$ will look like

$$0 \subseteq X_1 = N_1 \subseteq X_2 = N_1 \oplus N_2 \subseteq X_3 = N_1 \oplus N_2 \oplus N_3 \subseteq \cdots$$

where the N_i 's are direct sums of distinct indecomposables in $\mathfrak{s}$. Then $\cup_{i \in I} X_i = \bigoplus_{i \in I} N_i$ is an upper bound for the chain in $\mathcal{Z}$, so by Zorn's Lemma, $\mathcal{Z}$ has a maximal element, say X' , which is a pure submodule of X . Since X' is a direct sum of finitely many isomorphism types of modules of finite endolength, X' itself has finite endolength, and hence is pure injective [7, Chapter 4]. Since X' is a pure injective pure submodule, X' is a summand of X . So we have $X = X' \oplus X''$ with X'' having no non-zero finite length summand. By Corollary 5, X'' must be 0, and so $X = X'$ is a direct sum of finitely generated indecomposable submodules. $\square$

3. APPLICATION TO SUBSPACE REPRESENTATIONS OF POSETS

In this section we will let Λ be a right Morita ring. Let P be a poset, P^* the quiver obtained from P by adding a largest point, ΛP^* the incidence algebra of P , which is a free Λ -module with basis $\{(i, j) : i \leq j \text{ in } P^*\}$ as discussed in [2, Chapter 1]. Consider the full subcategory $\mathcal{S} = \text{Rep}_\Lambda P$, of $\text{Mod } \Lambda P^*$ which has as objects the subspace representations of P . First we will show that $\mathfrak{s} = \text{rep}_\Lambda P$ is functorially finite in $\text{mod } \Lambda P^*$ with respect to $\mathcal{S}$. With this we are able to describe $\mathcal{S} = \text{Rep}_\Lambda P^*$ in all cases where $\mathfrak{s}$ is of finite type, as we will see in one example.

Familiarity with the case of the one point poset will be useful in the proof of the next lemma, so for the moment, let $P = \bullet$ and $X \in \text{mod } \Lambda P^*$. Then X consists of a triple $(X_1, X_*; X_\alpha)$, where X_1, X_* are Λ -modules and $X_\alpha : X_1 \rightarrow X_*$. Let $\bar{e} : \text{Ker } X_\alpha \rightarrow I$ be the injective envelope of $\text{Ker } X_\alpha$. Using the injective factoring property, we can lift $\bar{e}$ to a map $e : X_1 \rightarrow I$. In [10] Ringel and Schmidmeier introduced $\text{Mimo}(X) = (X_1, X_* \oplus I_1; (X_\alpha, e)^T)$ and showed that the canonical map $\pi : \text{Mimo}(X) \rightarrow X$, is a right approximation for X in $\mathfrak{s}$. By the construction, we see the factorization properties hold for test objects in $\mathcal{S}$. So for the one point poset, $\text{mod } \Lambda P^*$ has right approximations in

$\mathfrak{s}$ with respect to $\mathcal{S}$. In the proof of the next lemma, we will generalize this construction for arbitrary posets. Notice that the assumption that R is a right Morita ring is needed so that the injective envelope of a finite length module is again of finite length.

Lemma 7. *Let P be a poset. The subcategory $\mathfrak{s} = \text{rep}_\Lambda P$ is functorially finite in $\text{mod } \Lambda P^*$ with respect to $\mathcal{S}$.*

Proof. For ease of notation, we label the vertices of P by natural numbers $1, 2, \dots, n$ via some ordering of the points which extends from the partial ordering. The notation $i < j$ will refer to the ordering of the poset. Let $X \in \text{mod } \Lambda P^*$. Then $X = (X_i, X_\alpha)$ where X_i is a Λ -module at vertex i , and X_α is a map $X_\alpha : X_{s(\alpha)} \rightarrow X_{t(\alpha)}$. For $i \leq j$ in P , and p a path from i to j , let X_p be the composition of the corresponding maps X_α . Since the map X_p is independent of the chosen path by the commutativity relations in $\text{mod } \Lambda P^*$, so we can refer to this morphism as X_{ij} .

With this set up, first we will show that $X \in \text{mod } \Lambda P^*$ has a left approximation in $\mathfrak{s}$ with respect to $\mathcal{S}$. Let $L(X) = (L_i, L_\alpha)$ where $L_i = \text{Im } X_{i*}$ and each $L_\alpha : L_{s(\alpha)} \rightarrow L_{t(\alpha)}$ is the inclusion map. Then $L(X)$ together with $l : X \rightarrow L(X)$ where $l = (X_{1*}, X_{2*}, \dots, X_{n*}, 1_{X*})$

fulfills the factorization property, and hence is a left approximation of X in $\mathfrak{s}$ with respect to $\mathcal{S}$.

It remains to show that X has a right approximation in $\mathfrak{s}$ with respect to $\mathcal{S}$. For this, we generalize the construction of Mimo recalled above. First, we define $\text{Mimo}_k(X)$ for a vertex k in P . Let $(I_k, \overline{e_k})$ be the injective envelope of the kernel of X_{k*} , and let $e_k : X_k \rightarrow I_k$ be the extension of $\overline{e_k}$ via the injective factoring property. Then $\text{Mimo}_k(X) = (M_i, M_\alpha)$ where

$$M_i = \begin{cases} X_i & i \leq k \\ X_i \oplus I_k & i \not\leq k \end{cases}$$

and for an arrow $\alpha : i \rightarrow j$ in P :

$$M_\alpha = \begin{cases} X_\alpha & j \leq k \\ (X_\alpha, e_k \circ X_{ik})^T & i \leq k, j \not\leq k \\ X_\alpha \oplus 1_{I_k} & i, j \not\leq k \end{cases}$$

So for each k , $\text{Mimo}_k(X)$ is a representation with a monomorphism starting at vertex k provided all arrows starting at successors of k are monomorphisms in the representation X . Hence

$$R(X) = \text{Mimo}_1(\text{Mimo}_2(\cdots (\text{Mimo}_n(X)) \cdots))$$

has monomorphisms for each arrow, so $R(X) \in \mathfrak{s}$. The module $R(X)$ together with the projection $r : R(X) \rightarrow X$ satisfies the factorization property required, since each map in the chain does:


$$R(X) = \text{Mimo}_1(\text{Mimo}_2(\cdots(\text{Mimo}_n(X))\cdots)) \rightarrow \cdots \rightarrow \text{Mimo}_n(X) \rightarrow X$$

Hence $r : R(X) \rightarrow X$ is a right approximation of X in $\mathfrak{s}$ with respect to $\mathcal{S}$. $\square$

As an example, let k be any field, $\Lambda = \frac{k[T]}{T^2}$, and P the poset with corresponding quiver:



Then $X \in \text{mod } \Lambda P^*$ is given by the following information:




To give a clear picture of the Mimo construction, below is the first step in computing $R(X)$:

$$\text{Mimo}_3(X) : \begin{array}{ccccc} & & X_1 & & \\ & \swarrow^{(e_3 \circ X_\beta)} & & \searrow^{X_\beta} & \\ I_3 \oplus X_2 & & & & X_3 \\ & \searrow^{1_{I_3} \oplus X_\gamma} & & \swarrow^{(e_3)} & \\ & & I_3 \oplus X_* & & \end{array}$$

And our right and left approximations $R(X)$ and $L(X)$ are:

$$L(X) : \begin{array}{ccc} & \text{Im}(X_\gamma \circ X_\alpha) & \\ \swarrow \iota & & \searrow \iota \\ \text{Im } X_\gamma & & \text{Im } X_\delta \\ \searrow \iota & & \swarrow \iota \\ & X_* & \end{array} \quad R(X) : \begin{array}{ccc} & X_1 & \\ \swarrow^{(e_3 \circ X_\beta)} & & \searrow^{(X_\beta)} \\ I_3 \oplus X_2 \oplus I_1 & & X_3 \oplus I_2 \oplus I_1 \\ \searrow^{1_{I_3} \oplus (X_\gamma)} & & \swarrow^{(e_3)} \\ I_3 \oplus X_4 \oplus I_2 \oplus I_1 & & \end{array}$$

Notice that $\mathcal{S} = \text{Rep}_\Lambda P$ is really the category of embeddings of embeddings of Λ -modules, and that the corresponding category of maps of maps, $\text{Mod } \Lambda P^*$ is known to be of infinite type. We compute a component Γ of the Auslander-Reiten quiver in $\mathcal{S}$ using coverings, approximations (as above), and methods from [11]. For a representation $X = (X_i)_{i \in P^*}$ we use the columns of boxes to denote the $\frac{k[T]}{T^2}$ module X_* corresponding to the total space at $(*)$, $\smile$ the image of a generator of the submodule X_2 , $\smile$ the image of a generator of X_3 , and $\bullet$ the image of a generator of X_1 .



For example, if

$$N = \begin{array}{c} \text{---} \text{---} \text{---} \\ \text{---} \text{---} \text{---} \end{array} \quad M = \begin{array}{c} \text{---} \text{---} \text{---} \\ \text{---} \text{---} \end{array}$$

Then M stands for the representation with $M_1 = M_2 = M_3 = M_* = \Lambda$ while N stands for the representation with $N_* = \Lambda \oplus \Lambda$, $N_1 = k \oplus 0$, $N_2 = k \oplus \Lambda$, and $N_3 = \Lambda \cdot (1, 1) \oplus k$ where k denotes the socle of Λ .

We can read off immediately that $\mathfrak{s}$ is of finite type since only 25 objects occur and Γ contains a projective generator. Since Λ is an artin algebra, each indecomposable is endofinite, and $\text{mod } \Lambda P^*$ has

Auslander-Reiten sequences in $\text{Mod } \Lambda P^*$. By Lemma 7 $\mathfrak{s}$ is functorially finite and hence has Auslander-Reiten sequences in $\mathcal{S}$ by proposition 4. Thus the conditions of Proposition 2 are satisfied, and for this example each object in $\mathcal{S}$ is a direct sum of finite length indecomposable modules. In other words, any indecomposable embedding of embeddings of $\frac{k[T]}{T^2}$ -modules has finite length. Also, as a consequence of Proposition 2, we obtain the following detailed information about the indecomposable summands of an arbitrary element of $\mathcal{S}$.

Corollary 8. *Let $X \in \text{Rep}_\Lambda P$ be an arbitrary representation. Then X has a direct sum decomposition into indecomposables such that each indecomposable summand Y occurs in our list of 25 indecomposables and has the following properties:*

- Y_* is a direct sum of at most two $\frac{k[T]}{T^2}$ -modules.
- Y is either isomorphic to N or both Y_2 and Y_3 are cyclic as $k[T]$ modules.
- Y is either isomorphic to M or $\dim Y_1 \leq 1$.

The above result can also be stated in terms of invariant subspaces of linear operators since $\frac{k[T]}{T^2}$ -modules are pairs (V, T) where V is a vector space and $T : V \rightarrow V$ is a linear operator acting with nilpotency index

2. Viewed in this way, submodules of $\frac{k[T]}{T^2}$ -modules are invariant subspaces, and we see that Proposition 3 is just a translation of Corollary 7 into the language of invariant subspaces.

4. PROOF OF THE MAIN RESULT

For this section, we require that R is an artinian ring, $\mathcal{S}$ contains a projective generator and is closed under direct sums and subobjects such that $\mathfrak{s} = \text{mod } R \cap \mathcal{S}$ is of finite type. Lemmas 9–14 leading up to a restatement of the Auslander and Ringel-Tachikawa Theorem have been copied or adapted from lecture notes of Zimmermann [14] where the classical version of the Auslander and Ringel-Tachikawa Theorem is shown. For the convenience of the reader, proofs are recalled.

Let $M_1, \dots, M_n$ be a list of all indecomposables in $\mathfrak{s}$ up to isomorphism, and let $M = M_1 \oplus \dots \oplus M_n$. Fix $X \in \mathcal{S}$, and let $H = \text{Hom}_R(M, X)$. We will denote $\text{Hom}_R(Y, Z)$ by (Y, Z) . Let $S = \text{End } M_R$. Then M is a left S -module, denoted ${}_S M$, and $(M, X)_S$ is a right S -module via the structure $g \circ s(m) = g(s(m))$.

Lemma 9. $X = \sum_{h \in H} \text{Im } h$

Proof. Let X' be a finitely generated submodule of X . The category $\mathcal{S}$ is closed under submodules, so $X' \in \mathcal{S}$ and hence in $\mathfrak{s}$. Since $\mathfrak{s}$ is of finite type, there are $k_1, \dots, k_n \in \mathbf{N}_0$ such that $X' \cong M_1^{k_1} \oplus \dots \oplus M_n^{k_n}$. So $X' \subseteq \sum_{h \in H} \text{Im } h$ and therefore $X = \sum_{h \in H} \text{Im } h$ $\square$

By this lemma, there is an epimorphism $p : M^{(H)} \rightarrow X$ given by $(m_h)_{h \in H} \mapsto \sum_{h \in H} h(m_h)$. With $K = \text{Ker } p$, we have a short exact sequence

$$\mathcal{E} : 0 \rightarrow K \rightarrow M^{(H)} \xrightarrow{p} X \rightarrow 0$$

Lemma 10. *The sequence $\mathcal{E} : 0 \rightarrow K \rightarrow M^{(H)} \xrightarrow{p} X \rightarrow 0$ is pure exact.*

Proof. Let $F \in \mathcal{S}$ be finitely presented. We need to check that the sequence

$$0 \rightarrow (F, K) \rightarrow (F, M^{(H)}) \xrightarrow{(F, p)} (F, X) \rightarrow 0$$

is exact, in particular, that (F, p) is an epimorphism. Let $f \in (F, X)$. Since F is finitely generated, F is a summand of M^m for some $m \in \mathbf{N}_0$. So there is a split monomorphism $j : F \rightarrow M^m$ and a morphism $q :$

$M^m \rightarrow F$ such that $1_F = q \circ j$. Let $\iota_i : M \rightarrow M^m$ be the i^{th} inclusion map. Then we have the following diagram.

$$\begin{array}{ccc}
 & & M \\
 & & \downarrow \iota_i \\
 & & M^m \\
 & & \uparrow j \quad \downarrow q \\
 & & F \\
 & & \downarrow f \\
 M^{(H)} & \xrightarrow{p} & X
 \end{array}$$

Let $f_i = f \circ q \circ \iota_i : M \rightarrow X$. Then $f_i \in H$ is the restriction of p to the summand of $M^{(H)}$ with index f_i , by definition of p . Denote by ι_{f_i} the canonical inclusion $\iota_{f_i} : M \rightarrow M^{(H)}$ into the f_i th component in $M^{(H)}$. Let $g : M^m \rightarrow M^{(H)}$ be the map induced by the ι_{f_i} so that $g \circ \iota_i = \iota_{f_i}$. Finally, let $h = g \circ j : F \rightarrow M^{(H)}$. Then we have the following commutative diagram.

$$\begin{array}{ccccc}
 & & M & & \\
 & & \downarrow \iota_i & & \\
 & & M^m & & \\
 & & \uparrow j \quad \downarrow q & & \\
 & & F & & \\
 & & \downarrow f & & \\
 M^{(H)} & \xrightarrow{p} & X & & \\
 \uparrow \iota_{f_i} & & \uparrow g & & \uparrow h \\
 M & & M^m & & F
 \end{array}$$

(Note: The diagram above is a simplified representation of the commutative diagram. The original diagram shows a more complex structure with curved arrows and labels f_i and h .)

Since $f_i = fq\iota_i = p\iota_{f_i}$, we have $pg\iota_i = p\iota_{f_i} = f_i = fq\iota_i$, for all i . So $pg = fq$, and $ph = pgj = fqj = f$. So for $f \in (F, K)$, there is an $h \in (F, M^{(H)})$ with $ph = f$; and (F, p) is onto. $\square$

Lemma 11. *The sequence obtained by applying $(M, -)$ to $\mathcal{E}$,*

$$0 \rightarrow (M, K) \rightarrow (M, M^{(H)}) \rightarrow (M, X) \rightarrow 0,$$

is a pure exact sequence of S -modules.

Proof. Since $0 \rightarrow K \rightarrow M^{(H)} \rightarrow X \rightarrow 0$ is a pure exact sequence of R -modules,

$$0 \rightarrow (M, K) \rightarrow (M, M^{(H)}) \rightarrow (M, X) \rightarrow 0,$$

is an exact sequence of S -modules by Lemma 10 since M is finitely presented. It remains to show that the sequence is pure exact.

Let N be a finitely generated right S -module. Then there is an epimorphism $\pi : S^n \rightarrow N$ for some n . Since $-\otimes_S M : \text{Mod } S \rightarrow \text{Mod } R$ is a right exact functor, we get an R epimorphism $\pi \otimes 1 : S^n \otimes_S M \rightarrow N \otimes_S M$. Since tensor products commute with direct sums, $S^n \otimes_S M = M^n$ is a finitely generated R -module. Hence $N \otimes_S M_R = \text{Im}(\pi \otimes 1)$ is also a finitely generated R -module.

Now we are ready to apply $(N, -)$ to our sequence and check that it gives a short exact sequence.

$$0 \rightarrow (N_S, (M_R, K_R)_S) \rightarrow (N_S, (M_R, M_R^{(H)})_S) \rightarrow (N_S, (M_R, X_R)_S)$$

Notice that for a right S -module N , an S - R bimodule M , and a right R -module Y , we have the adjoint isomorphism which is natural in N and Y :

$$(N \otimes_S M, Y) \rightarrow (N, \text{Hom}_R(M, Y))$$

$$\phi \mapsto (n \mapsto (m \mapsto \phi(n \otimes m)))$$

So we obtain the commutative diagram:

$$\begin{array}{ccccccc} 0 & \longrightarrow & (N_S, (M_R, K_R)_S) & \longrightarrow & (N_S, (M_R, M_R^{(H)})_S) & \longrightarrow & (N_S, (M_R, X_R)_S) \\ & & \uparrow \cong & & \uparrow \cong & & \uparrow \cong \\ 0 & \longrightarrow & (N_S \otimes M_R, K_R) & \longrightarrow & (N_S \otimes M_R, M_R^{(H)}) & \longrightarrow & (N_S \otimes M_R, X_R) \end{array}$$

But $0 \rightarrow K \rightarrow M^{(H)} \rightarrow X \rightarrow 0$ is pure exact and $N_S \otimes M_R$ is finitely generated, so the bottom sequence is short exact. Hence $(N_S, (M_R, M_R^{(H)})_S) \rightarrow (N_S, (M_R, X_R)_S)$ is an epimorphism by the commutativity of the diagram; and so the lemma is proved. $\square$

Lemma 12. *The evaluation map $\lambda : (M, X) \otimes_S M \rightarrow X_R$ where $\phi \otimes m \mapsto \phi(m)$ is an isomorphism of R -modules.*

Proof. Let $x \in X_R$. Then $x = \sum \phi_i(m_i)$ for some $\phi_i \in (M, X)$ and $m_i \in M$ by Lemma 9. It remains to show that λ is injective. Let $\sum_{i=1}^m \phi_i \otimes x_i \in \text{Ker } \lambda$, with $\phi_i \in (M, X)$, $x_i \in M$. Then $\lambda(\sum_{i=1}^m \phi_i \otimes x_i) = \sum_{i=1}^m \phi_i(x_i) = 0$. Also, since $R_R \in \mathcal{S}$, Lemma 9 implies we can write $1 = \sum_{j=1}^n \alpha_j(m_j)$ for some $\alpha_j \in (M, R)$, $m_j \in M$. Let $s_{ij} : M \rightarrow M$ be the R -homomorphism given by $m \mapsto x_i \alpha_j(m)$. Then

$$\begin{aligned}
 (1) \quad \sum_{i=1}^m \phi_i \otimes x_i &= \sum_{i=1}^m \phi_i \otimes x_i \cdot 1 \\
 (2) \quad &= \sum_{i=1}^m \phi_i \otimes x_i \sum_{j=1}^n \alpha_j(m_j) \\
 (3) \quad &= \sum_{i,j} \phi_i \otimes s_{ij}(m_j) \\
 (4) \quad &= \sum_{i,j} \phi_i s_{ij} \otimes m_j \\
 (5) \quad &= \sum_j \left(\sum_i \phi_i s_{ij} \right) \otimes m_j \\
 (6) \quad &= \sum_j 0 \otimes m_j \\
 (7) \quad &= 0
 \end{aligned}$$

Note that equality (4) holds since $s_{ij} \in S$, and we can see that (6) holds since for $x \in M$, $\sum_i \phi_i s_{ij}(x) = \sum_i \phi_i(x_i \alpha_j(x))$ but $\alpha_j(x)$ is a scalar and $\sum_{i=1}^m \phi_i(x_i) = 0$ by the kernel assumption. So $\text{Ker } \lambda = 0$; and we see that λ is a monomorphism and hence an isomorphism, as required. $\square$

Lemma 13. $(M, M^{(H)}) \cong S^{(H)}$ as right S -modules.

Proof. Since $S = \text{End}(M)$, $S^{(H)} = \bigoplus_H(M, M)$. Also, $(M, M^{(H)}) = (M, \bigoplus_H M)$. But M is finitely generated, so $\bigoplus_H(M, M) = (M, \bigoplus_H M)$. $\square$

In particular, since $S^{(H)}$ is a projective S -module; $(M, M^{(H)}) \cong S^{(H)}$ is a projective S -module. To prove the main result, however, we would like (M, X) to be projective. Since S is the endomorphism ring of a finite length module, it is semiprimary; and so we will use the following general lemma.

Lemma 14. *Let S be semiprimary. Let*

$$\mathcal{E} : 0 \rightarrow A \xrightarrow{f} B \xrightarrow{g} C \rightarrow 0$$

be a pure exact sequence of S -modules with B projective. Then C is also projective.

Proof. Since S is semiprimary, every S -module has a projective cover, and each projective in S is the direct sum of indecomposable projective S -modules. [1, Theorem 27.11] Let $\pi : P \rightarrow C$ be a projective cover of C , $P = \bigoplus_{i \in I} P_i$ for some set I and indecomposable projective S -modules P_i . We want to show that $L = \text{Ker}(\pi)$ is zero, so pick $x \in L$. Since $x \in L \subseteq P = \bigoplus_{i \in I} P_i$, there is a finitely generated projective summand P' of P , with $x \in P'$. Since $\pi : P \rightarrow C$ is an epimorphism and $g : B \rightarrow C$ with B projective, the projective factoring property yields a map $t : B \rightarrow P$ such that $\pi t = g$. So for the kernel map t' , we have the following commutative diagram with exact rows.

$$\begin{array}{ccccccccc}
 0 & \longrightarrow & A & \xrightarrow{f} & B & \xrightarrow{g} & C & \longrightarrow & 0 \\
 & & \downarrow t' & & \downarrow t & & \parallel & & \\
 0 & \longrightarrow & L & \xrightarrow{\iota} & P & \xrightarrow{\pi} & C & \longrightarrow & 0
 \end{array}$$

Let $j : P' \rightarrow P$ be the inclusion map. Since $x \in P'$, xS is a submodule of P' , so let $\iota' : xS \rightarrow P'$ be this inclusion map. Then we have

$$\begin{array}{ccccccccc}
 0 & \longrightarrow & A & \xrightarrow{f} & B & \xrightarrow{g} & C & \longrightarrow & 0 \\
 & & \downarrow t' & & \downarrow t & & \parallel & & \\
 0 & \longrightarrow & L & \xrightarrow{\iota} & P & \xrightarrow{\pi} & C & \longrightarrow & 0 \\
 & & & & \uparrow j & & & & \\
 0 & \longrightarrow & xS & \xrightarrow{\iota'} & P' & \xrightarrow{\pi'} & \frac{P'}{xS} & \longrightarrow & 0
 \end{array}$$

Since $x \in L$, $xS \subseteq L$, and there is an inclusion map $j' : xS \rightarrow L$, which makes the bottom left square commute and hence induces a cokernel map $j'' : \frac{P'}{xS} \rightarrow C$. Also, P' is a summand of P , and j is a split monomorphism, so there is a q such that $1_{P'} = qj$.

$$\begin{array}{ccccccccc}
 0 & \longrightarrow & A & \xrightarrow{f} & B & \xrightarrow{g} & C & \longrightarrow & 0 \\
 & & \downarrow t' & & \downarrow t & & \parallel & & \\
 0 & \longrightarrow & L & \xrightarrow{\iota} & P & \xrightarrow{\pi} & C & \longrightarrow & 0 \\
 & & \uparrow j' & & \downarrow q & & \uparrow j & & \\
 0 & \longrightarrow & xS & \xrightarrow{\iota'} & P' & \xrightarrow{\pi'} & \frac{P'}{xS} & \longrightarrow & 0
 \end{array}$$

But $\frac{P'}{xS}$ has projective resolution $P_1 \rightarrow P' \xrightarrow{\pi'} \frac{P'}{xS}$, where P_1 is finitely generated since it is the projective cover of $\text{Ker } \pi' = xS$. Since $\frac{P'}{xS}$ is also finitely generated, it is finitely presented. Also, $\mathcal{E}$ is pure exact, so for $j'' \in \text{Hom}(\frac{P'}{xS}, C)$ there is a $\tau \in \text{Hom}(\frac{P'}{xS}, B)$ such that $j'' = g\tau$. Since $\pi(j - t\tau\pi') = j''\pi' - j''\pi' = 0$ by commutativity, $j - t\tau\pi' \in \text{Ker } \pi = \text{Im } \iota$, and there is a $\sigma : P' \rightarrow L$ so that $j - t\tau\pi' = \iota\sigma$. But $\iota\sigma\iota' = (j - t\tau\pi')\iota' = j\iota' - t\tau\pi'\iota' = j\iota' = \iota j'$ by commutativity and

exactness of the third row, and $\iota\sigma\iota' = \iota j'$ implies $\sigma\iota' = j'$ since ι is injective.

Now let $h = \iota\sigma q \in \text{End } P$. Since $\pi h = \pi\iota\sigma q = 0$, $\text{Im } h \subseteq \text{Ker } \pi$. But π is a projective cover, so $\text{Ker } \pi$ is small in P , and hence $\text{Im } h$ is also small in P . So $h \in \text{Rad}(\text{End}(P))$. Since S is semiprimary; $\text{Rad}(\text{End}(P))$ is nilpotent; so h is nilpotent, and $1 - h$ is an automorphism of P . Finally, $(1 - h)(j\iota'(x)) = j\iota'(x) - \iota\sigma q j\iota'(x) = j\iota'(x) - \iota\sigma\iota'(x) = 0$. Since $1 - h$ is an automorphism; $j\iota'(x) = 0$. Since $j\iota'$ is a monomorphism; $x = 0$, so $L = 0$. Hence $P \cong C$ and we see C is projective. $\square$

Since M is a finitely generated right R module in $\mathcal{S}$, and $S = \text{End}(M_R)$ we have the functor

$$(- \otimes_S M_R) : \mathbf{P}_S \rightarrow \mathbf{S}_M$$

where $\mathbf{P}_S$ is the category of projective right S -modules and $\mathbf{S}_M$ is the category of direct summands of direct sums of copies of M [1, Lemma 29.4]. We can now complete the proof of Theorem 1:

Proof. Let $X \in \mathcal{S}$. Let $M_1, \dots, M_n$ be a list of all indecomposables in $\mathcal{S}$ up to isomorphy, define $M = M_1 \oplus \dots \oplus M_n$, and $S = \text{End}(M)$. Then

the short exact sequence

$$0 \rightarrow (M, K) \rightarrow (M, M^{(H)}) \rightarrow (M, X) \rightarrow 0$$

is pure exact by Lemma 11. By Lemma 13, the middle term is projective, and Lemma 14 yields that $(M, X)_S$ is projective. Applying the functor $(- \otimes_S M_R)$ to $(M, X)_S \in \mathbf{P}_S$, we see $(M, X)_S \otimes_S M_R \in \mathbf{S}_M$. By Lemma 12, $(M, X)_S \otimes_S M_R \cong X$, so $X \in \mathbf{S}_M$ and hence is a direct summand of $M^{(J)}$ for some set J . By definition of M , X is a direct summand of $M_1^{(J)} \oplus \cdots \oplus M_n^{(J)}$. So by Azumaya's theorem; $X = M_1^{(J_1)} \oplus \cdots \oplus M_n^{(J_n)}$ where $J_1, \cdots, J_n \subseteq J$. $\square$

REFERENCES

- [1] F.W. Anderson and K.R. Fuller, *Rings and categories of modules*, Springer-Verlag, GTM **13** (1992).
- [2] I. Assem, D. Simson, and A. Skowroński, *Elements of the representation theory of associative algebras*, London Mathematical Society, ST **65** (2006).
- [3] M. Auslander, *Functors and morphisms determined by objects*, Rep. Theory of Algebras: Proc. Phil. Conf., NY (1978), 1–244.
- [4] M. Auslander, *Representation theory of artin algebras II*, Comm. Alg. **1** (1974), 269–310.
- [5] M. Auslander and S. Smalø, *Almost split sequences in subcategories*, J. Alg. **69** (1981), 126–151.

- [6] M. Auslander, I. Reiten, and S. Smalø, *Representation theory of artin algebras*, Cambridge Press, **36** (1997).
- [7] W. Crawley-Boevey, *Modules of finite length over their endomorphism rings*, Cambridge University Press, LNS **168** (1992), 127–184.
- [8] F. Richman and E. Walker, *Subgroups of p^5 -bounded subgroups*, Abelian groups and modules, Birkhäuser, (1999), 55–73.
- [9] C.M. Ringel, *Foundation of the representation theory of artin algebras, using the Gabriel-Roiter measure*, Contemporary Mathematics, **406** (2006), 105–135.
- [10] C.M. Ringel and M. Schmidmeier, *The Auslander-Reiten translation in submodules categories*, Trans. Amer. Math. Soc., **360** (2008), 691–716.
- [11] C.M. Ringel and J. Schröer, *Representation theory of algebras I*, Second Preliminary Version.
- [12] C.M. Ringel and H. Tachikawa, *QF3 rings*, J. Reine Angew. Math. **272** (1975), 49–72.
- [13] W. Zimmermann, *Auslander-Reiten sequences over artinian rings*, J. Alg. **119** (1988), 366–392.
- [14] W. Zimmermann, *Einführung in die Darstellungstheorie*, Lecture notes, winter semester 1980-1981, Universität München (LMU).